

LA-UR-20-24076

Approved for public release; distribution is unlimited.

Title: Quarkonium: experimental overview

Author(s): Durham, J. Matthew

Intended for: Hard Probes 2020, 2020-06-01 (Austin, Texas, United States)

Issued: 2020-06-04





Outline



Introduction

- $Q\bar{Q}$ in Small Systems \rightarrow Large Systems
 - Focus on medium interactions

- Exotic hadrons containing $Q\bar{Q}$
 - X(3872) in the QCD medium
- Brief look to the future

Outline




• Introduction

- $Q\bar{Q}$ in Small Systems \rightarrow Large Systems
 - Focus on medium interactions

- Exotic hadrons containing $Q\bar{Q}$
 - X(3872) in the QCD medium
- Brief look to the future

Heavy $Q\overline{Q}$ states in vacuum






Solve Schrodinger equation with the potential

$$V_0^{(c\bar{c})}(r) = -\frac{4}{3}\frac{\alpha_s}{r} + br + \frac{32\pi\alpha_s}{9m_c^2}\tilde{\delta}_{\sigma}(r)\vec{S}_c \cdot \vec{S}_{\bar{c}}$$

Phys. Rev. D 72, 054026 (2005)

Heavy $Q\bar{Q}$ states in vacuum






Solve Schrodinger equation with the potential


$$V_0^{(c\bar{c})}(r) = -\frac{4}{3}\frac{\alpha_s}{r} + br + \frac{32\pi\alpha_s}{9m_c^2}\tilde{\delta}_{\sigma}(r)\vec{S}_c \cdot \vec{S}_{\bar{c}}$$


Phys. Rev. D 72, 054026 (2005)



Heavy $Q\bar{Q}$ states in vacuum








Measured mass: $3842.71 \pm 0.16 \pm 0.12 \text{ MeV}$

Predicted mass: 3849 MeV

Incredibly rich structure, accessible theoretically and experimentally




Heavy $Q\bar{Q}$ states in the QCD medium



Diffuse medium

Increasing T, N_{ch}


Dense medium



Dissociation via interactions with comoving particles


Heavy $Q\bar{Q}$ states in the QCD medium






Increasing T, N_{ch}

Dense medium






Hydrodynamic flow induced by pressure gradients (initial state?)


Heavy $Q\bar{Q}$ states in the QCD medium







Increasing T, N_{ch}

Dense medium






Hydrodynamic flow induced by pressure gradients (initial state?)




Heavy QQ states in the QCD medium





Heavy $Q\overline{Q}$ states in the QCD medium





^{*}Additional initial state effects from modified nPDF and energy loss can also be significant in collisions with nuclei.

Experimentally, we use different collision systems/kinematic regions to prepare environments where different effects dominate for different probes.

Every experiment contributes to quarkonia physics




Taken together:

$$-5 < y < 5$$


0< p_T <several hundred GeV














Outline




Introduction

- $Q\bar{Q}$ in Small Systems \rightarrow Large Systems
 - Focus on medium interactions

- Exotic hadrons containing $Q\bar{Q}$
 - X(3872) in the QCD medium
- Brief look to the future

Charmonium in pp vs. multiplicity







New ALICE results on J/ψ production in 13 TeV pp shows faster-than-linear growth with multiplicity: various CGC calculations match data

Charmonium in pp vs. multiplicity






New ALICE results on J/ψ production in 13 TeV pp shows faster-than-linear growth with multiplicity: various CGC calculations match data




Hints that $\psi(2S)$ production does now grow as fast as J/ψ with multiplicity – initial state models should give similar trend for both. Final state effect suppressing $\psi(2S)$ in pp?

Charmonium in pp vs. multiplicity







New ALICE results on J/ψ production in 13 TeV pp shows faster-than-linear growth with multiplicity: various CGC calculations match data



Hints that $\psi(2S)$ production does now grow as fast as J/ψ with multiplicity – initial state models should give similar trend for both. Final state effect suppressing $\psi(2S)$ in pp?









Midrapidity STAR data shows little J/ψ modification at intermediate p_T :

- Consistent with nPDF calculations
- Consistent with PHENIX d+Au data
- CNM not dominant effect in AA in this range






Forward rapidity (p or 3He-going direction):

• Data generally in agreement with nPDF calculations Backward rapidity (A-going direction):

- Data from Au nucleus is **inconsistent with nPDF calculation**:
 - Additional effects required to explain data

Significantly improved precision in state-of-the art nPDF calculations, which include D, J/ψ , and Y(1S) LHC data ion:

Kusina, Lansberg, Schienbein, Shao, Phys. Rev. Lett. 121, 052004 (2018)



Forward rapidity (p or 3He-going direction):

• Data generally in agreement with nPDF calculations Backward rapidity (A-going direction):


- Data from Au nucleus is **inconsistent with nPDF calculation**:
 - Additional effects required to explain data


Significantly improved precision in state-of-the art nPDF calculations, which include D, J/ψ , and $\Upsilon(1S)$ LHC data ion:


Kusina, Lansberg, Schienbein, Shao, Phys. Rev. Lett. 121, 052004 (2018)

$Q\bar{Q}$ in pPb









At both forward and backward rapidity, J/ψ well described by nPDF calculations: initial state dominates


- Marked difference from PHENIX data, where additional effects are required to match backwards J/ψ data
- J/ ψ breakup smaller at LHC smaller crossing time


Backwards $\psi(2S)$ data shows inconsistency with nPDF


Florian Damas, Mon 1:35

$Q\bar{Q}$ in pPb









Florian Damas, Mon 1:35

At both forward and backward rapidity, J/ψ well described by nPDF calculations: initial state dominates

- Marked difference from PHENIX data, where additional effects are required to match backwards J/ψ data
- J/ ψ breakup smaller at LHC smaller crossing time


Backwards $\psi(2S)$ data shows inconsistency with nPDF


• Tension eases when final state comover breakup is included, small effect for forward.


Similar effects previously observed by PHENIX, ALICE, LHCb Similar sequential suppression of Υ (nS) also observed in pPb

$Q\bar{Q}$ in pPb









Florian Damas, Mon 1:35

At both forward and backward rapidity, J/ψ well described by nPDF calculations: initial state dominates

- Marked difference from PHENIX data, where additional effects are required to match backwards J/ψ data
- J/ ψ breakup smaller at LHC smaller crossing time


Backwards $\psi(2S)$ data shows inconsistency with nPDF

Tension eases when final state comover breakup is included, small effect for forward.

Similar effects previously observed by PHENIX, ALICE, LHCb Similar sequential suppression of Υ (nS) also observed in pPb





Data on J/ ψ production is consistent with nPDF, however: Significant J/ ψ v₂ observed in pPb collisions





Data on J/ ψ production is consistent with nPDF, however: Significant J/ ψ v₂ observed in pPb collisions







• Similar to PbPb; some differences observed at high pt, typically attributed to L dependent energy loss, not present in pPb



Data on J/ ψ production is consistent with nPDF, however: Significant J/ ψ v₂ observed in pPb collisions







- Similar to PbPb; some differences observed at high pt, typically attributed to L dependent energy loss, not present in pPb
- Transport model with short lived plasma phase in pPb does not give large v2 in pPb (does describe low pT PbPb)



Data on J/ ψ production is consistent with nPDF, however: Significant J/ ψ v₂ observed in pPb collisions






- Similar to PbPb; some differences observed at high pt, typically attributed to L dependent energy loss, not present in pPb
- Transport model with short lived plasma phase in pPb does not give large v2 in pPb (does describe low pT PbPb)
- CGC calculation describes part of pPb v2; however CGC has mixed results describing light sector v2

MAJOR OPEN QUESTION




Data on J/ ψ production is consistent with nPDF, however: Significant J/ ψ v₂ observed in pPb collisions




• CGC calculation gives very similar v2 for Upsilon states on pPb: next challenge for experiments

$\Upsilon(nS)$ in PbPb






Textbook example of color screening




- Three well defined state observable in $\mu^+\mu^-$
- Bottom density relatively low expect minimal effects from recombination

$\Upsilon(nS)$ in PbPb






Textbook example of color screening Well described by models



- Three well defined state observable in $\mu^+\mu^-$
- Bottom density relatively low expect minimal effects from recombination


$\Upsilon(nS)$ in PbPb



Textbook example of color screening Well described by models









Generally good agreement between ATLAS and CMS

$Q\bar{Q}$ flow in PbPb







In contrast to J/ψ , no significant v_2 is observed for Y(1S) by ALICE or CMS. Little or no recombination due to relative low b density in QGP, as expected from models.

Outline



Introduction

- $Q\bar{Q}$ in Small Systems \rightarrow Large Systems
 - Focus on medium interactions

- Exotic hadrons containing $Q\bar{Q}$
 - X(3872) in the QCD medium


Brief look to the future


Hard Probes 1-9: 0 talks devoted to X(3872)

Hard Probes 10: 4 talks! 2 experiment, 2 theory

Exotic $Q\overline{Q}$ states: XYZ









Exotic $Q\bar{Q}$ states: XYZ






Dozens of states containing $c\bar{c}$ and $b\bar{b}$ have been discovered that are not predicted by any $Q\bar{Q}$ potential model

Compact tetraquark




Hadronic molecule




Thousands of theory papers exploring possible structures, but no consensus on even the first exotic, X(3872)

Primarily measured as products of b decays in vacuum




Exotic $Q\bar{Q}$ states: XYZ




Dozens of states containing $c\bar{c}$ and $b\bar{b}$ have been discovered that are not predicted by any $Q\bar{Q}$ potential model

Compact tetraquark


Hadronic molecule





Interactions in QCD medium are dependent on binding energy/structure


Production of multiquark exotics is especially sensitive to recombination/coalescence effects that are expected in heavy ion collisions




Exotic X(3872) in diffuse medium (8 TeV pp)

Los Alamos NATIONAL LABORATORY EST 1943

LHCb-CONF-2019-005



- Prompt X(3872)/ ψ (2S) decreases with activity
 - X(3872) suppressed more than $\psi(2S)$




Systematic uncertainty box fully correlated point-to-point

Cameron Dean, Wed 12:25

Exotic X(3872) in diffuse medium (8 TeV pp)


Los Alamo

LHCb-CONF-2019-005



- Prompt X(3872)/ ψ (2S) decreases with activity
 - X(3872) suppressed more than $\psi(2S)$
- *B*-decay component consistent with constant
 - Production in vacuum unaffected

Cameron Dean, Wed 12:25




Systematic uncertainty box fully correlated point-to-point

Exotic X(3872) in diffuse medium (8 TeV pp) Los A



LHCb-CONF-2019-005




- Prompt X(3872)/ ψ (2S) decreases with activity
 - X(3872) suppressed more than $\psi(2S)$
- B-decay component consistent with constant
 - Production in vacuum unaffected

Cameron Dean, Wed 12:25

$$\sigma^{\text{co}-\mathcal{Q}}(E^{\text{co}}) = \sigma_{\text{geo}}^{\mathcal{Q}} \times \left(1 - \frac{E_{\text{thr}}^{\mathcal{Q}}}{E^{\text{co}}}\right)^n$$

Directly probes size σ_{geom} and binding energy





tetraquark interpretation

Elena Ferreiro, Wed 1:05

Exotic X(3872) in dense medium (PbPb)








Recombination of X(3872) at pT > 15 GeV?

Prompt X(3872)/ ψ (2S) = 1.10 ± 0.51 ± 0.53 in PbPb at 5 TeV Prompt X(3872)/ ψ (2S) \approx 0.1 in pp at 8 TeV

Exotic X(3872) in dense medium (PbPb)





Prompt X(3872)/ ψ (2S) \approx 0.1 in pp at 8 TeV

Intriguing data! Inconclusive with these uncertainties.

Outline




Introduction

- $Q\bar{Q}$ in Small Systems \rightarrow Large Systems
 - Focus on medium interactions

- Exotic hadrons containing $Q\bar{Q}$
 - X(3872) in the QCD medium
- Brief look to the future

Fixed target at LHCb: SMOG II





https://cds.cern.ch/record/2673690/files/LHCB-TDR-020.pdf

Upgraded SMOG system at LHCb allows greatly increased rates of beam+injected gas collisions

No centrality limitations in p+gas or Pb+gas at LHCb

Barbara Trzeciak (AFTER physics program), Wed 1:55

Samuel Belin (Future of LHCb), Thurs 10:55


Example SMOG2 pAr at 115 GeV for one year

Int. Lumi.	80 pb ⁻¹
Sys.error of J/Ψ xsect J/Ψ yield D^0 yield Λ_c yield Ψ' yield $\Upsilon(1S)$ yield $DY \mu^+\mu^-$ yield	ion ~3% 28 M 280 M 2.8 M 280 k 280 k 24 k 24 k


Large heavy flavor samples
Access to exotic states near RHIC energies

New detector at RHIC: sPHENIX





Rosi Reed, Fri 8:00




Precision bottomonium measurements at RHIC are a major focus of sPHENIX

Future facility: electron-ion collider

E. Aschenauer, Wed 9:15






EIC site selection at BNL announced Jan 2020

Charm production inside the nucleus probes:

- Parton structure of nucleons
- nPDF modifications
- QCD energy loss

Xuan Li, Thurs 10:35

In the kinematic range accessed by the EIC, hadronization *inside the nucleus* becomes an important effect on observables



Future facility: electron-ion collider




EIC site selection at BNL announced Jan 2020

Charm production inside the nucleus probes:

- Parton structure of nucleons
- nPDF modifications
- QCD energy loss

Xuan Li, Thurs 10:35

e+A collisions have relatively low backgrounds – well suited for studies of higher charmonia and exotics: P_c^+ , Z_c^+ , χ_c , ...



Summary



- Quarkonia remains a major focus of heavy ion physics 34 years after Matsui and Satz
 - Contributions from all experimental collaborations a wealth of data
 - Strong synergy with theory community
- The lines between small and large systems are continually becoming blurred
 - Final state suppression manifests in many systems at high multiplicity
 - Universal understanding of J/ ψ flow from small to large systems remains elusive
- Exotic multiquark states are becoming accessible in a QCD medium
 - Medium probes exotic properties \Leftrightarrow Exotics probe medium properties
- Exciting future programs of in-medium quarkonia studies at LHC/RHIC/EIC to come




Los Alamos is supported by the US Dept. of Energy/Office of Science/Office of Nuclear Physics




BACKUPS

We have only measured 2 charmonium states in heavy ion collisions at LHC.



We have only measured 3 bottomonium states in heavy ion collisions at LHC.





Quark Model of Hadrons



Volume 8, number 3

PHYSICS LETTERS

1 February 1964

AN SU_{z} MODEL FOR STRONG INTERACTION SYMMETRY AND ITS BREAKING

A SCHEMATIC MODEL OF BARYONS AND MESONS *

M. GELL-MANN

California Institute of Technology, Pasadena, California

Received 4 January 1964

A simpler and more elegant scheme can be constructed if we allow non-integral values for the charges. We can dispense entirely with the basic baryon b if we assign to the triplet t the following properties: spin $\frac{1}{2}$, $z=-\frac{1}{3}$, and baryon number $\frac{1}{3}$. We then refer to the members u^3 , $d^{-\frac{1}{3}}$, and $s^{-\frac{1}{3}}$ of the triplet as "quarks" 6) q and the members of the anti-triplet as anti-quarks \bar{q} . Baryons can now be constructed from quarks by using the combinations (qqq), $(qqq\bar{q})$, etc., while mesons are made out of $(q\bar{q})$, $(qq\bar{q}\bar{q})$, etc. It is assuming that the lowest baryon configuration (qqq) gives just the representations 1, 8, and 10 that have been observed, while the lowest meson configuration $(q\bar{q})$ similarly gives just 1 and 8.

G.Zweig *)
CERN - Geneva


8182/TH.401 17 January 1964

In general, we would expect that baryons are built not only from the product of three aces, AAA, but also from AAAAA, AAAAAAAA, etc., where A denotes an anti-ace. Similarly, mesons could be formed from AA, AAAA etc. For the low mass mesons and baryons we will assume the simplest possibilities, AA and AAA, that is, "deuces and treys".


States with >3 quarks have been expected since the beginning of the quark model

Charmonium in pp - RHIC

PHENIX: PRD 101 (2020) 5, 052006



Agreement with calculations across entire p_T range





STAR, *PRD* 100 (2019) 5, 052009


Charmonium in pp - LHC



ATLAS-CONF-2019-047

Preliminary result from ATLAS:

 J/ψ measured up to 360 GeV $\psi(2S)$ measured up to 140 GeV



Charmonium in pp - LHC





ATLAS-CONF-2019-047


Preliminary result from ATLAS:

 J/ψ measured up to 360 GeV $\psi(2S)$ measured up to 140 GeV

Non-prompt component (from b decays) diverges from FONLL calculations for $p_T > 100 \text{ GeV}$










J/ψ flow: pPb and PbPb













Improved nPDF extraction








This data is already being used to constrain the gluon nPDF down to x~5x10⁻⁶


Kusina, Lansberg, Schienbein, Shao, Gluon shadowing and antishadowing in heavy-flavor production at the LHC Phys. Rev. Lett. 121, 052004 (2018)




Phys. Lett. B Volume 774, 10 November 2017, Pages 159-178




$Q\bar{Q}$ in pPb












At both forward and backward rapidity, J/ ψ and Y(1S) states are well described by nPDF calculations: initial state dominates

• Marked difference from PHENIX data, where additional effects beyond nPDF were required to match backwards data

$Q\bar{Q}$ in pPb







At both forward and backward rapidity, J/ψ and $\Upsilon(1S)$ states are well described by nPDF calculations: initial state dominates Final state effects required to explain suppression of excited states at backwards rapidity



\overline{D}^* Molecule


state	η_c	J/ψ	χ_{c0}	χ_{c1}	χ_{c2}	ψ'	X(3872)
mass [GeV]	2.98	3.10	3.42	3.51	3.56	3.69	3.872
$\Delta E \text{ [GeV]}$	0.75	0.64	0.32	0.22	0.18	0.05	$\begin{array}{c} 0.00001 \pm \\ 0.00027 \end{array}$

Satz, J. Phys. G 32 (3) 2006

Exotic X(3872) in diffuse medium (8 TeV pp)

• Los Alamos NATIONAL LABORATORY EST. 1943


LHCb-CONF-2019-005



Prompt fraction of X(3872) and ψ (2S) both drop as function of event activity:


- Increasing *b* component?
- Suppressed prompt component? i.e. final state effect in *pp*?

Cameron Dean, Wed 12:25



"No significant contribution from coalescence" at pT > 10 GeV/c for charm baryons






Quarkonia inside the nucleus




Fixed target p+A experiments also operated in a kinematic region where quarkonia hadronized inside the nucleus – similar to EIC




Significant differences observed between states with same quark content, but different binding energies


Natural explanation based on size: Larger (weakly bound) states sample a larger volume of the nucleus while passing through – larger absorption cross section

Arleo, Gossiaux, Gousset, Aichelin PRC 61 (2000) 054906










Recent LHCb Discovery – P_c States




Select daughters from the decay

$$\Lambda_b^0 \to J/\psi p K^-$$

Masses are close to meson+baryon thresholds – candidate for hadronic molecule







Charged Exotic – Z_c State



Select daughters from the decay





Charged and contains $c\bar{c}$: Minimal 4 quark content: $c\bar{c}q\bar{q}$

Mass is NOT close to any hadron+hadron threshold – candidate for compact tetraquark